

**USING STABLE ISOTOPE PROBING AND BIO-TRAPS TO IDENTIFY
TBA DEGRADERS *IN SITU* UNDER SULFATE REDUCING CONDITIONS**

Kerry Sublette*

Center for Applied Biogeosciences
The University of Tulsa
800 South Tucker Drive
Tulsa, OK 74104
Voice: 918-631-3085
Fax: 918-631-3268
kerry-sublette@utulsa.edu

Jennifer Busch-Harris

ConocoPhillips
Bartlesville, OK

Kathleen Duncan

University of Oklahoma
Norman, OK

Greg Davis

Dora Ogles

Microbial Insights, Inc.
Rockford, TN

Doug Mackay

University of California-Davis
Davis, CA

We have demonstrated that biofilms characteristic of aquifer conditions can be rapidly and efficiently collected using *in situ* microcosms or “bio-traps” containing Bio-Sep® beads. Bio-Sep® beads are 3-4 mm in diameter and composed of 25% aramid polymer and 75% powdered activated carbon (PAC). They have a porous, sponge-like structure with 74% porosity and 600 m²/g of surface area surrounded by an ultrafiltration-like membrane that contains holes and tears of 1-10 microns in size. Bio-Sep® beads are heated to 300 °C for sterilization and to render the beads free of fossil biomarkers.

When bio-traps are deployed in groundwater, indigenous bacteria enter through the outside membrane and migrate into the porous internal matrix. Microbes then attach to this internal structure and reproduce to form biofilms. In this manner, microbes can be concentrated for analysis despite their relatively low density of microbes in the sampled groundwater. Microorganisms must grow and reproduce within the bead to be detected. Thus the beads will collect only those organisms which are active under the specific subsurface conditions. Those microorganisms that are not capable of growth under *in situ* conditions due to a nutrient limitation, unfavorable redox potential, or adverse environmental conditions will not be collected in the beads at detectable levels.

Bio-Sep® beads can also be “baited” with a variety of organic compounds by vapor phase adsorption onto the PAC component of the beads. The adsorbed organic has been shown to be bioavailable to bacteria that form biofilms in the beads during incubation in a contaminated aquifer. If the compound is labeled with ¹³C, polar lipids may be extracted from bead biofilms and the derived fatty acid methyl esters analyzed for ¹³C incorporation using GC-IRMS. Since the beads are clean of biomarkers and sterile when deployed, incorporation of ¹³C in phospholipids provides proof of *in situ* biodegradation of a target compound by indigenous microorganisms under actual aquifer conditions.

We have successfully used bio-traps amended with ¹³C₆-benzene, ¹³C₅-MTBE, and ¹³C₄-TBA to demonstrate the *in situ* biodegradation potential of these compounds at several sites. One such site has been the Vandenberg AFB site where bio-traps amended with ¹³C₄-TBA were deployed in an anaerobic TBA plume. Several phospholipid fatty acids extracted from bead biofilms were found to be labelled with ¹³C indicating biodegradation of TBA. Fatty acid and DNA labelling patterns suggest the participation of sulfate-reducing bacteria in the degradation process.

###